


**AMENDMENTS TO THE CLAIMS**


1. (previously presented) A diaminobenzene compound represented by formula (1) below



where  $R^1$  and  $R^2$  each independently denotes a hydrogen atom, alkyl group, or alkoxy group.

2. (previously presented) The diaminobenzene compound as defined in claim 1, wherein  $R^1$  and  $R^2$  each independently denotes a  $C_{1-20}$  alkyl group,  $C_{1-20}$  alkoxy group, or  $C_{1-20}$  fluoroalkyl group.


3. (previously presented) A polyimide precursor which comprises repeating units represented by formula (2) below



where  $R^1$  and  $R^2$  each independently denotes a hydrogen atom, alkyl group, or alkoxy group;


"A" denotes a residue of tetracarboxylic acid; and n denotes an integer of 1 to 5000.

4. (previously presented) A polyimide which comprises repeating units represented by formula (3) below



where R<sup>1</sup> and R<sup>2</sup> each independently denotes a hydrogen atom, alkyl group, or alkoxy group;  
 "A" denotes a residue of tetracarboxylic acid; and n denotes an integer of 1 to 5000.

5. (currently amended) A polyimide precursor which is obtained by reaction between  
 a diamine component containing at least 1 mol% of ~~the diaminobenzene compound~~  
~~defined in claim 1 or 2~~ a diaminobenzene compound represented by formula (1) below




where R<sup>1</sup> and R<sup>2</sup> each independently denotes a hydrogen atom, alkyl group, or alkoxy group and

a tetracarboxylic acid or a derivative thereof.

6. (original) The polyimide precursor as defined in claim 5, wherein the tetracarboxylic acid or the derivative thereof is an aromatic tetracarboxylic acid or a derivative thereof.

7. (original) The polyimide precursor as defined in claim 6, wherein the aromatic tetracarboxylic acid is a tetracarboxylic acid having phenyl groups or substituted phenyl groups.

8. (currently amended) A polyimide which is obtained by ring-closing reaction from any of polyimide precursors as defined in claim 5 obtained by reaction between a diamine component containing at least 1 mol% of a diaminobenzene compound represented by formula (1) below



where R<sup>1</sup> and R<sup>2</sup> each independently denotes a hydrogen atom, alkyl group, or alkoxy group and a tetracarboxylic acid or a derivative thereof.

9. (previously presented) A charge carrier transporting film which is formed from the polyimide as defined in claim 4.

10. (previously presented) An organic transistor device which comprises the charge carrier transporting film as defined in claim 9.

11. (original) An organic light emitting diode which has at least one layer of the charge carrier transporting film as defined in claim 9.

12. (previously presented) A fluorescent filter which comprises the charge carrier transporting film as defined in claim 9.

13. (previously presented) A liquid crystal alignment film which comprises the charge carrier transporting film as defined in claim 9.

14. (new) The polyimide precursor as defined in claim 5, wherein  $R^1$  and  $R^2$  each independently denotes a  $C_{1-20}$  alkyl group,  $C_{1-20}$  alkoxy group, or  $C_{1-20}$  fluoroalkyl group.

15. (new) The polyimide as defined in claim 8, wherein  $R^1$  and  $R^2$  each independently denotes a  $C_{1-20}$  alkyl group,  $C_{1-20}$  alkoxy group, or  $C_{1-20}$  fluoroalkyl group.